Physica Scripta

INCLUSIVE PUBLISHING TRUSTED SCIENCE

PURPOSE-LED PUBLISHING™

PAPER • **OPEN ACCESS**

Alena Tensor in unification applications

To cite this article: Piotr Ogonowski and Piotr Skindzier 2025 *Phys. Scr.* **100** 015018

View the article online for updates and enhancements.

You may also like

- Tensor Quantum Programming
Alena A. Termanova, Artem A. Melnikov, Egor Mamenchikov et al.
- VIPRE: A Tool Aiding the Design for Entry Probe Missions
Alena Probst, Linda Spilker, Mark Hofstadter et al.
- Flare Expansion to a Magnetic Rope Accompanied by Rare Radio Bursts
Alena Zemanová, Marian Karlický, Jana Kašparová et al.

This content was downloaded from IP address 178.205.54.240 on 16/12/2024 at 15:10

Physica Scripta

PAPER

Alena Tensor in unification applications

Piotr Ogonowski[1] **and Piotr Skindzier**[2]

[1] Kozminski University, Jagiellonska 57/59, Warsaw, 03-301, Poland
[2] Jagiellonian University, Alumni, Reymonta 4, Krakow, 30-059, Poland

E-mail: piotrogonowski@kozminski.edu.pl

Keywords: Unification, Electrodynamics, General Relativity, Quantum Field Theory, Quantum Mechanics, Dark Matter, Dark Energy

OPEN ACCESS

RECEIVED
6 September 2024

REVISED
13 November 2024

ACCEPTED FOR PUBLICATION
29 November 2024

PUBLISHED
10 December 2024

Original content from this work may be used under the terms of the Creative Commons Attribution 4.0 licence.

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Abstract

Alena Tensor is a recently discovered class of energy-momentum tensors that provides mathematical framework in which, as demonstrated in previous publications, the description of a physical system in curved spacetime and its description in flat spacetime with fields are equivalent. The description of a system with electromagnetic field based on Alena Tensor can be used to reconcile physical descriptions. (1) In curvilinear description, Einstein Field equations were obtained with Cosmological Constant related to the invariant of the electromagnetic field tensor, which can be interpreted as negative pressure of vacuum, filled with electromagnetic field. (2) In classical description for flat spacetime, three densities of four-forces were obtained: electromagnetic, against gravity (counter-action to gravitational free-fall), and the force responsible for the Abraham-Lorentz effect (radiation reaction force). Obtained connection of Einstein tensor with gravity and radiation reaction force, after transition to curvilinear description, excludes black hole singularities. There was obtained Lagrangian density and generalized canonical four-momentum, containing electromagnetic four-potential and a term responsible for the other two forces. In this description charged particles cannot remain at complete rest and should have spin, their energy results from the existence of energy of magnetic moment and the density of this energy is part of the Poynting four-vector. The distribution of charged matter was expressed as polarization-magnetization stress-energy tensor, what may explain why gravity is invisible in QED. 3) In quantum picture, QED Lagrangian density simplification was obtained, and the Dirac, Schrödinger and Klein–Gordon equations may be considered as approximations of the obtained quantum solution. Farther use of Alena Tensor in unification applications was also discussed.

1. Introduction

The history of physics is also the history of unification. The past teaches us that after the stage of research on individual phenomena and obtaining a satisfactory description of them, comes the phase of unification, in which the scattered puzzles of knowledge are put together into one whole picture, which soon turns out to also be just a part of bigger picture.

Today, modern physicists are faced with many puzzles, most of which are huge pictures, entire sections of physics, composed of hundreds of smaller parts, the existence of which we owe to thousands of outstanding scientists. The largest and most famous descriptions of physical phenomena requiring unification are, of course, General Relativity (GR) and Quantum Field Theory (QFT), however, the unification cannot be simplified to finding a theory of quantum gravity. We cannot forget about other knowledge components (so fundamental that they are easy to miss), such as Continuum Mechanics or Thermodynamics, which are also being researched in the field of unification [1–3]. It is also important to note that even after a quantum picture of gravity is obtained, questions about the Dark Sector will still remain open [4].

'In all the attempts at unification we encounter two distinct methodological approaches: a deductive-hypothetical and an empirical-inductive method' [5]. where a good examples of the first approach are String Theory [6] and Supersymmetry [7] and of the second one, Grand Unification [8] and, in a sense, the Standard

© 2024 The Author(s). Published by IOP Publishing Ltd

Model itself. Part of the entire unification effort are dualistic theories [9], mainly adopting mentioned deductive-hypothetical approach. They are usually looking for a theoretical model in which existing descriptions can be reconciled and assume, that contradictions between existing descriptions may be apparent and in fact they are only different, equally valid ways of describing the same phenomena [10].

Considering the context of unification broadly, a dualistic solution to the puzzle may appear from a completely unexpected direction, as in the work of D. Grimmer describing topological redescription [11] and giving the possibility of changing the topology of space in a way similar to changing coordinate systems. When considering the unification of GR and Electrodynamics, a dualistic unification theory may come from a rather obvious direction, connecting the description of electrodynamics with the geometry of spacetime [12, 13], because it can be expected that there is a mathematical transformation between accelerated motion in flat spacetime and geodesic motion in curved spacetime for all accelerations due to considered fields.

This paper will analyze a new unifying dualistic approach, called the Alena Tensor, using it to describe a physical system with an electromagnetic field. Previous publications [14, 15] have already shown, that Alena Tensor allows to obtain a coherent solution combining relativistic electrodynamics, continuum mechanics, QED and GR equations, so it is not just a purely theoretical, mathematical construction and seems worth further development. This method also indicates that the description of the physical system in curved spacetime and its description in flat spacetime with fields are equivalent, thanks to an appropriately constructed definition of the energy-momentum tensor for the system, which allows analysis in both flat and curved spacetime. Thanks to the equivalence of descriptions, as will be shown in this article, in the resulting picture of interactions in flat spacetime, gravity is not a force, which is indeed a direct consequence of GR. The force related to gravity occurring in a system in flat spacetime, will be found to act against gravity (e.g. centrifugal force) and result from any motion other than free fall, respecting equivalence principle.

In Alena Tensor approach, the metric tensor is not a property of spacetime, but only a way of describing it. The tensor of the field present in the system is responsible for all forces in flat spacetime, and thus also the field tensor defines the metric tensor for which all forces vanish and may be completely replaced by curvature in curvilinear description. According to this approach, gravity is not a body force (in the sense of Continuum Mechanics), but a side effect of existence of the energy of other, more fundamental fields (e.g. EM, Electroweak). As was shown in [14, 15], creation of the energy-momentum tensor taking into account the above assumptions, allows for a dualistic description (curvilinear, classical and quantum), leads to compatibility with currently used descriptions of physical systems and opens the way to many further studies. In this paper this analysis will be continued, using smooth transitions between the description of gravity, electromagnetism and the quantum picture, which is possible with the use of the Alena Tensor.

Another and perhaps the most important reason to write this article is that the Alena Tensor is not an intuitive theory, requires some systematization and yet requires further research. Therefore, to facilitate analysis and future development, all main conclusions from previous publications have been systematized and aggregated in the appendix. Since this article adds another piece of knowledge to the already existing conclusions, such a summary seems necessary. Otherwise, all the extensive reasoning from the previous articles would have to be presented in this chapter.

Appendix focuses on how the Alena Tensor can be used to solve the unification puzzle and as a new approach, it may seem counter-intuitive. Analysis of unconventional reasoning (even if it has already been analyzed, discussed with peer reviewers and published) can be difficult and discouraging for the reader, but if unification did not require new approach, we would have known about it long ago. For instance, description of a physical theory usually begins with a description of the action and by varying it, one finds the equations of the theory, energy-momentum tensor and Lagrangians. In this case, however, such a line of reasoning would make it difficult to understand the unifying potential of this theory, which is why the action and the Lagrangian (derived in previously published papers) appear only later in the appendix.

When considering a curved spacetime, metrics are typically obtained from the solutions of the GR equations based on the symmetries used. In this article, the conclusions regarding the Einstein tensor will be presented mainly in flat spacetime, which also breaks a certain accepted pattern and is not intuitive. However, such an analysis will reveal the meaning of the presented dualistic description without the need to significantly expand this article, especially since Alena Tensor by definition transforms into Einstein Field Equations in curvilinear description and the methods of analyzing GR equations are quite well known. The article will therefore focus on flat spacetime, showing that the equivalence principle is satisfied in the Alena Tensor framework, explaining the method for calculating the free-fall velocity and for calculating the force necessary for the existence of stationary observers above the gravity source. The general solution for a dynamical system with gravity in flat spacetime will be also discussed, and it will be explained, how to reproduce in flat spacetime known static GR solutions using Alena Tensor.

In the Results section the authors will interpret and develop the results obtained in previous publications (described in appendix) for the system with electromagnetic field, showing that it leads to the conclusions

described in the abstract. The beginning of the Results section will introduce the reader to the notation and further reasoning to make the equations understandable also to the readers not familiar with Alena Tensor. The rest of the article will discuss the possibilities of further development and applications of Alena Tensor to analyze problems related to the broadly understood research on the unification of physical theories.

The authors use the Einstein summation convention, metric signature $(+,-,-,-)$ and commonly used notations.

2. Results

This chapter will consider the physical system with only electromagnetic field, in the way described by Alena Tensor according to appendix. However, since the considerations concern mainly flat spacetime, this simplifies most of the equations and for the analysis of this chapter it may be enough to know the main definitions for flat spacetime:

ϱ_o is rest mass density, γ is Lorentz gamma factor, and ϱ following reasoning related to equation (11) of [14] is consistently denoted as $\varrho \equiv \varrho_o\gamma$. Therefore, mass density in four-momentum density ϱU^α is equal to $\varrho_o\gamma^2 = \varrho\gamma$. In flat spacetime also (A3) and (A5) can be simplified to

$$\Lambda_\rho \equiv \frac{1}{4\mu_o}\mathbb{F}^{\alpha\beta}\mathbb{F}_{\alpha\beta}\ ;\ p \equiv \varrho c^2 + \Lambda_\rho \tag{1}$$

where $\mathbb{F}^{\alpha\beta}$ represents the electromagnetic field tensor, μ_o is vacuum magnetic permeability, Λ_ρ is negative and p is negative pressure in the system. The remaining definitions and equations will be introduced successively with references to the conclusions obtained so far, included in appendix.

According to the conclusion obtained (A17), three four-force densities appear in the considered system in flat spacetime: electromagnetic f_{EM}, related to gravity f_{gr} and other f_{oth}. Since the least obvious element is f_{oth}, it will be interpreted in section 2.1. This four-force density is negligibly small for $-\Lambda_\rho >> \varrho c^2$ and as it will be shown, it appears to reproduce Abraham–Lorentz effect [16] (also called the radiation reaction force). Section 2.1 will also discuss the four-force density f_{gr} and its relation to the interpretation of gravity in curved spacetime. A relativistic description of motion in flat spacetime reproducing the Schwarzschild metric will also be derived as an example.

It is also known, that the existence of magnetic moment is expected for charged particles [17, 18] and it influences the value of the electromagnetic force [19, 20]. It will be therefore shown in section 2.2, that the use of the energy-momentum tensor for the system in form of Alena Tensor (denoted as $T^{\alpha\beta}$) leads directly to the conclusion that charged particles should have spin and the energy of the particles results from the existence of energy of magnetic moment. For this reason, the energy of charged particles will also turn out to be part of the Poynting four-vector and the previously obtained relationship between four-momentum and derived gauge of electromagnetic four-potential (A24) will be clarified. The presence of the charged matter energy density in the Poynting four-vector can be already seen by analyzing the first row of the classical energy-momentum tensor of electromagnetic field $\Upsilon^{\alpha\beta}$ (A4) in flat spacetime, which using conclusion (A6) can be alternatively expressed as

$$\Upsilon^{0\beta} = \frac{\Lambda_\rho}{p}\varrho\ c\gamma U^\beta - \frac{\Lambda_\rho}{p}T^{0\beta} \tag{2}$$

In above Υ^{00} represents the energy density of electromagnetic field and from (A27) it is known, that $T^{00} = \Lambda_\rho$ thus $-\frac{\Lambda_\rho^2}{p}$ and $\frac{\Lambda_\rho}{p}\varrho\ c^2\gamma^2$ are positive, so the above equation represents the components of the Poynting four-vector. However, the connection of $\frac{\Lambda_\rho}{p}\varrho\ c^2\gamma^2$ with the energy density of magnetic moment requires more detailed explanation in section 2.2.

In section 2.3, the Alena Tensor verification method within QM will be presented and it will be demonstrated, that the Dirac, Schrödinger and Klein–Gordon equations may be considered as approximations of the obtained quantum solution, resulting from the use of the Alena Tensor. It will be also shown, that presented approach allows combining the classical and quantum descriptions of the motion of charged particles which can help in many applications.

2.1. Interpretation of the four-force densities in flat spacetime

Using (1), one may define relative permeability μ_r and volume magnetic susceptibility χ as

$$\mu_r \equiv \frac{\Lambda_\rho}{p}\ ;\ \chi \equiv \mu_r - 1 = -\frac{\varrho c^2}{p} \tag{3}$$

thanks to which the Alena Tensor (A6) in flat spacetime takes the form expected for the system with electromagnetic field

$$T^{\alpha\beta} = \varrho\ U^{\alpha}U^{\beta} - \frac{1}{\mu_r}\Upsilon^{\alpha\beta} \tag{4}$$

It now may be noticed, that there are present in the system (A17) only two four-force densities: related to gravity f^{α}_{gr} according to (A26) equal to

$$f^{\alpha}_{gr} = \Upsilon^{\alpha\beta}\partial_{\beta}\frac{1}{\mu_r} = \varrho\left(c^2\partial^{\alpha}\ln(\mu_r) - \frac{d\ \ln(\mu_r)}{d\tau}U^{\alpha}\right) \tag{5}$$

and electromagnetic corrected by the $\frac{1}{\mu_r}$ coefficient

$$f^{\alpha}_{EM} + f^{\alpha}_{oth} = \left(1 + \frac{\varrho c^2}{\Lambda_{\rho}}\right)\partial_{\beta}\Upsilon^{\alpha\beta} = \frac{1}{\mu_r}f^{\alpha}_{EM} \tag{6}$$

Introduced μ_r allows for the interpretation of the applied correction to the electromagnetic four-force density, linking the presence of relative permeability with the existence of energy density of matter. In the limit for $\varrho c^2 = 0$ Alena Tensor simply becomes a tensor of the electromagnetic field $-T^{\alpha\beta} = \Upsilon^{\alpha\beta}$ and pressure $p = \Lambda_{\rho}$. Therefore Λ_{ρ} may be actually associated with the negative pressure of a vacuum, filled with an electromagnetic field. In the limit $\varrho c^2 = -\Lambda_{\rho}$ one obtains $p = 0$ and forces caused by field disappear. The meaning of this boundary solution requires further analysis.

Using (A33), (A35), (A37) and definition of the pressure (A5) one may notice, that considering point-like particles one obtains

$$cW^0 = W_{pv} = -\int p\ d^3x\ ;\ H = -\int \Lambda_{\rho}\ d^3x = mc^2\gamma + W_{pv} \tag{7}$$

where H is Hamiltonian for point-like particle, W_{pv} is pressure-volume work and there is relation of the pressure p (A5) in the system to the energy of the field cW^0 contained in the system. It thus also becomes possible to interpret the correction for the electromagnetic force discussed in (6) considering point-like particles, where now relative permeability μ_r and derived gauge of electromagnetic four-potential $q\mathbb{A}^{\mu}$ (A24) are

$$\frac{1}{\mu_r} = \frac{W_{pv}}{H}\ ;\ -q\mathbb{A}^{\mu} = \mu_r\ P^{\mu} = \frac{\chi H}{c^2}(\ c\ ,\ \overrightarrow{u}) \tag{8}$$

As one may notice from (7) and (A37), the increasing energy of an accelerated body cannot take energy from nowhere. Considering H as conserved energy in a closed system, one gets that $mc^2\gamma$ increases at the expense of decreasing W_{pv} in the system. Therefore, also forces resulting from decreasing W_{pv} must at some point decrease and for $W_{pv} = 0$ the body cannot accelerate any more.

Considering the analyzed system as a medium in which the speed of light is equal to c and assuming classical relation between permeability $\mu = \mu_o\mu_r$ and permittivity $\varepsilon = \varepsilon_o\varepsilon_r$

$$\frac{1}{c^2} = \mu\cdot\varepsilon = \mu_o\varepsilon_o\cdot\mu_r\varepsilon_r \tag{9}$$

one also gets that relative permittivity ε_r and the electric susceptibility χ_e are

$$\varepsilon_r \equiv \frac{1}{\mu_r} = \frac{W_{pv}}{H}\ ;\ \chi_e \equiv \varepsilon_r - 1 = \frac{\varrho c^2}{\Lambda_{\rho}} = -\frac{mc^2\gamma}{H} \tag{10}$$

As may be seen, $f^{\alpha}_{oth} = \chi_e\, f^{\alpha}_{EM}$ acts as a negative correction to four-force densities as $mc^2\gamma$ increases at the expense of W_{pv} and upholds the principle of conservation of energy. It means, that discussed f_{oth} reproduces Abraham–Lorentz effect and represents radiation reaction force.

The Alena Tensor in curved spacetime transforms into EFE (A13) with the cosmological constant related to Λ_{ρ} (A14), (A3), but above conclusions allow already for the interpretation of f^{α}_{gr} in flat spacetime with a negligible electromagnetic force. The reader who uses GR equations in curved spacetime may be accustomed to using the stationary observer's coordinate time at infinity t_{∞}, metric for curved spacetime and the conserved energy E_o resulting from GR. However, since flat spacetime is considered in this section, in all the equations it is used $ct = X^0$ associated with the four-position X^{μ}, Lorentz factor $\gamma = \frac{dt}{d\tau}$ and the Minkowski metric is used, so the energy of the test body will be designated as E. In the following steps it will be shown that despite the use of flat spacetime, it is possible to reproduce the motion of a test body along a geodesic using the gravitational force f^{α}_{gr} resulting from Alena Tensor.

At first step, to improve the clarity of the equations and to hold the equation (10), one may introduce some dimensionless potential ϕ, such that

$$W_{pv} \equiv (H - E)e^{\phi - \frac{E_o}{mc^2}}\ \ \rightarrow\ \ \varepsilon_r = \frac{W_{pv}}{H} = 1 - \frac{mc^2\gamma}{H} \tag{11}$$

which links E_o from GR with E and ensures compliance with the equation (10), providing property

$$mc^2\phi = E_o \;\rightarrow\; \varepsilon_r = 1 - \frac{E}{H} \;\rightarrow\; mc^2\gamma = E \tag{12}$$

Above may be further simplified by assuming that the energy E_o conserved in GR is actually equal to $E_o = H - E$, however, this is not necessary for further analysis.

Thanks to (11) also (5) may be simplified to

$$\frac{1}{\varrho} f^{\alpha}_{gr} = \frac{d\phi}{d\tau} U^{\alpha} - c^2 \partial^{\alpha}\phi \tag{13}$$

According to the equivalence principle, it must be assumed that for free fall f^{α}_{gr} vanishes. Denoting the free fall velocity as $\overrightarrow{u_{ff}}$, equivalence principle therefore yields

$$-\overrightarrow{u_{ff}} = c\,\frac{\nabla\phi}{\partial^0\phi} \;\rightarrow\; \frac{d\phi}{dt} = \left(1 - \frac{\overrightarrow{u}\;\overrightarrow{u_{ff}}}{c^2}\right)\partial_t\phi \tag{14}$$

With the above condition, four-force density f^{α}_{gr} actually vanishes in free-fall ($\overrightarrow{u} = \overrightarrow{u_{ff}}$) and for $\partial^{\alpha}\phi = 0$.

Gravity is not a force what is direct consequence of GR. It drives to conclusion, that f^{α}_{gr} does not represent gravity itself, but the four-force density against gravity, resulting from the motion of a considered test body. f^{α}_{gr} must therefore take into account the test body's motion and the associated counteraction to gravity. In stationary, symmetric cases, for a test body in a fixed position from the source of gravity (stationary observer) with $U^{\alpha} \rightarrow (c, 0)$ four-force density f^{α}_{gr} should be therefore equal to gravitational force with opposite sign, representing the force needed to stay in fixed place. For circular orbits with $\frac{d\phi}{d\tau} = 0$ and $U^{\alpha} \neq (c, 0)$ it can be expected, that $\partial^{\alpha}\phi$ should vanish. Therefore ϕ should be related to the effective potential and for symmetric, stationary cases it must be approximated by the Newtonian potential.

One may therefore use known effective potential resulting from Schwarzschild metric [21] as illustration, assuming L to be normalized angular momentum

$$\phi \equiv \sqrt{\frac{E_o^2}{m^2c^4} - \left(\frac{1}{c}\frac{dr}{d\tau}\right)^2} = \sqrt{\left(1 - \frac{r_s}{r}\right)\left(1 + \frac{L^2}{r^2}\right)} \tag{15}$$

$$\partial_r\,\phi = \frac{(\alpha - \beta)(\alpha + \beta)}{2r\phi}\;;\quad \alpha \equiv \sqrt{\frac{r_s}{r}\left(1 + \frac{L^2}{r^2}\right)}\;;\quad \beta \equiv \sqrt{\frac{2L^2}{r^2}\left(1 - \frac{r_s}{r}\right)} \tag{16}$$

Thanks to the above approach, in the limit $\frac{dr}{d\tau} = 0$ one obtains consistency with (12) and all effective potentials resulting from GR may be successfully used

$$\frac{dr}{d\tau} = 0 \;\rightarrow\; mc^2\phi = E_o \tag{17}$$

In the limit of radial only motion in considered Schwarzschild case, L vanishes, thus

$$L = 0 \;\rightarrow\; \beta = 0 \;\rightarrow\; \alpha = \sqrt{\frac{r_s}{r}} \tag{18}$$

Therefore in order to agree (16) with (14) and to keep consistency with radial-only case, one may expect, that

$$\partial^0\phi = \frac{\alpha - \beta}{2r\phi}\;;\quad -\overrightarrow{u_{ff}} = (\alpha + \beta)\,\frac{\overrightarrow{r}}{r}\,c \tag{19}$$

thus in radial-only case

$$-\overrightarrow{u_{ff}} = c\,\sqrt{\frac{r_s}{r}}\left(\frac{\overrightarrow{r}}{r}\right) \;\rightarrow\; \partial^0\phi = \frac{\sqrt{\frac{r_s}{r}}}{2r\phi} \tag{20}$$

Above may be interpreted as stationary observers being accelerated by time-invariant gravitational acceleration a_g, passed by freely falling body moving only in radial axis with time-invariant radial velocity u_{ff} opposite to theirs, with some displacement denoted as r_s where

$$r_s \equiv \frac{a_g\,t^2}{2}\;;\quad -u_{ff} \equiv a_g\,t\;;\quad a_g = -\partial_t\,u_{ff} = \partial_t^2\,r_s \tag{21}$$

In the above description, the Schwarzschild radius corresponds to the total path traveled by, in fact, stationary observers counteracting gravity. However, this is only a side effect of the assumptions made, where r_s is actually the path 'not traveled' by a radially falling body in free fall towards the point $r = 0$. It is also worth noting that the considered above dependence of r_s on t is possible (t is four-position time), where in general, dependence of r_s on



Figure 1. Coefficient values corresponding to the Schwarzschild case.

t_∞ is also essentially necessary to allow for the existence of Hawking radiation in Schwarzschild case [22]. Without such dependence, the mass of the gravitational source would remain constant, which would prevent any energy from being radiated.

For cases $L \neq 0$ for circular orbits the radius of the rotation r_{rot} is

$$\partial^\alpha \phi = 0 \quad \rightarrow \quad r_{rot} = \frac{L^2 \pm \sqrt{L^4 - 3L^2 r_s^2}}{r_s} \quad \rightarrow \quad L = \frac{r_{rot}\ \sqrt{r_s}}{\sqrt{2r_{rot} - 3r_s}} \tag{22}$$

what is also correct result for considered case. Obtained in this way f_{gr}^α with ϕ defined in (15) is an exact, flat spacetime description reproducing Schwarzschild metric preserving all the properties of the curvilinear description and shedding new light on the interpretation of gravity described by this metric. For example:

- four-force density f_{gr}^α vanishes only for $\alpha = \beta$ and $\overrightarrow{u} = \overrightarrow{u_{ff}}$
- case $\alpha = \beta$ represents circular orbit and in this case $\partial^0 \phi = 0$ what indicates that r_s and L does not change in time
- in all other cases L must be time dependent, since $\partial^0 \phi \neq 0$ which means that the body lose angular momentum as it moves around the source of gravity
- the energy lost in this way must be radiated in the form of gravitational waves, which de facto agrees with the general knowledge about GR
- bodies on orbits with $\partial^0 \phi \neq 0$ must in fact slowly spiral toward the source of gravity

By transforming the GR description into the flat spacetime description, the relativistic four-force density f_{gr}^α visible in (13) takes into account effective potential ϕ and the four-velocity U_α to provide property $U_\alpha f_{gr}^\alpha = 0$, reflect the complexity of the results obtained using GR and retain the conclusion that gravity is not a force, while preserving equivalence principle. Therefore, obtained from Alena Tensor relativistic four-force density f_{gr}^α, opens a new way to test different ϕ distributions describing gravity in flat spacetime, which may also help in the analysis of GR solutions, extending present knowledge.

The reasoning presented in this section can be used to reproduce gravity in flat spacetime also for gravity described by other metrics in curved spacetime. It can also be noticed that for radial free-fall with $E_o = mc^2$, even below the Schwarzschild radius, coefficients ε_r and μ_r have analyzable imaginary values, representing certain wave functions. This can be seen in figure 1 below, where ε_r and μ_r have been divided/multiplied by a constant $1 - \frac{E}{H}$ (normalized).

The above preliminary conclusions require further research and in-depth analysis for f_{gr}^α in flat spacetime corresponding to other metrics in curvilinear description. However, these conclusions are sufficient to make an analysis of the equation (4) from the perspective of electromagnetism, because, as one may notice, the energy-momentum tensor for the electromagnetic field $\Upsilon^{\alpha\beta}$ with μ_r (related to ϕ) defined in a similar way as above for other types of gravity sources, has also analyzable values both above and below the event horizon. This analysis will be performed in the next section.

Finally, it is worth noting that since four-force density associated with the Einstein tensor (accurate to a constant) in (A19), may be expressed in flat spacetime as

$$\partial_\beta G^{\alpha\beta} = f_{gr}^\alpha + f_{oth}^\alpha = \partial_\beta \chi_e \Upsilon^{\alpha\beta} \tag{23}$$

therefore f^{α}_{oth}, maintaining the principle of conservation of energy, must prevent non-physical effects such as black hole singularity. This can be seen by analyzing (in a classical way used for GR), solutions of (A13) in curved spacetime for the static, symmetric case, as these are smooth de Sitter solutions [23], free of singularities. However, this topic deserves to be developed in a separate article.

The interpretation presented in this section, introduces also new possibilities regarding the interpretation of the dark sector, which will be discussed later in the article.

2.2. Interpretation of the system from the perspective of electromagnetism theory

Staying with the description for a flat spacetime with an electromagnetic field, denoting the electric and magnetic fields as $\overrightarrow{E}$ and $\overrightarrow{B}$ and denoting the densities of electric and magnetic energy occurring in the electromagnetic field tensor as

$$u_{E\sim} \equiv \varepsilon_o \frac{E^2}{2}\ ;\ u_{B\sim} \equiv \frac{1}{\mu_o}\frac{B^2}{2} \tag{24}$$

it can be seen from (A3) that

$$\Lambda_\rho = u_{B\sim} - u_{E\sim} \tag{25}$$

Therefore using conclusion (A30), the relationship between magnetic energy density and the energy density of the electromagnetic field Υ^{00} can be written as

$$\frac{B^2}{\mu_o} = \Lambda_\rho + \Upsilon^{00} = \frac{\Lambda_\rho}{p}\varrho_o c^2(\gamma^3+\gamma) \tag{26}$$

Thanks to the above, the four-potential of the electromagnetic field from conclusion (A24) can be simplified to

$$\rho_o \mathbb{A}^\mu = -\frac{B^2}{\mu_o(\gamma^2+1)}\frac{1}{c^2\gamma}U^\mu \tag{27}$$

For a charged particle at rest, the above reduces to a scalar $-\frac{1}{c}\frac{B^2}{2\mu_o}$ expressing (negative) classical value of magnetic energy density and zero vector, but completely stationary cases must be excluded, because they lead to $\overrightarrow{B} = \nabla \times \overrightarrow{\mathbb{A}} = 0$. The above equation thus also says, that even in the absence of orbital angular momentum, the charged particle must vibrate or rotate and experience a magnetic field, because without the magnetic field, the entire four-potential vanishes. In fact, it will be shown later in this section, that the above four-potential results from the existence of magnetic moment and magnetization itself is the source of the electric field of quasi-stationary charged particles.

Using four-current $J^\alpha = \rho_o \gamma U^\alpha$ from (A29), the source of the electric field associated with charged matter can now be represented, as reduced (compared to the classical value) magnetic energy density $u_{B\odot}$

$$u_{B\odot} \equiv \frac{1}{\mu_o}\frac{B^2}{(\gamma^2+1)} = -J_\mu \mathbb{A}^\mu = \frac{\Lambda_\rho\, \varrho c^2}{p} = -\chi\Lambda_\rho \tag{28}$$

where χ is volume magnetic susceptibility from (3). In the above description, the $1+\gamma^2$ coefficient seems to be related to some intrinsic, internal volume magnetic susceptibility of charged matter, so one may take a closer look at this phenomenon. The value preserved in the system (A27) must be also conserved for the electromagnetic energy densities associated with charged matter

$$-\Lambda_\rho = u_{E\sim} - u_{B\sim} = u_{E\odot} - u_{B\odot} \tag{29}$$

where according to (28), (A5) and above, electric energy density associated with matter $u_{E\odot}$ may be denoted as

$$u_{E\odot} \equiv -\frac{\Lambda_\rho^2}{p} = -\mu_r\Lambda_\rho \tag{30}$$

The above leads directly to the conclusion that total electromagnetic field energy density may be expressed as electric field energy density related to charged matter and the energy density of magnetic moment. It may be seen by calculating energy density of the electromagnetic field

$$\Upsilon^{00} = u_{E\sim} + u_{B\sim} = u_{E\odot} + 2u_{B\sim} - u_{B\odot} = u_{E\odot} + \frac{\gamma^2}{(\gamma^2+1)}\frac{B^2}{\mu_o} \tag{31}$$

In above, last component of the equation represents the classical description of the energy density of magnetic moment, where γ^2 serves as volume magnetic susceptibility. Therefore, the electromagnetic field associated with density of charged matter will be most easily described as a propagating disturbance of magnetization and polarization, because the combination of magnetization and polarization describes such electric currents [24] and relativistic tensor can be created based on them [25]. Recalling the classical refuels of the theory of electromagnetism, by decomposing electromagnetic field tensor $\mathbb{F}^{\alpha\beta}$ into Polarization-Magnetization tensor

denoted e.g. as $\mathcal{M}^{\alpha\beta}$ and Electric Displacement tensor denoted e.g. as $\mathcal{D}^{\alpha\beta}$ one obtains $\frac{1}{\mu_o}\mathbb{F}^{\alpha\beta} = \mathcal{M}^{\alpha\beta} + \mathcal{D}^{\alpha\beta}$ where $\mathcal{M}^{\alpha\beta}$ and $\mathcal{D}^{\alpha\beta}$ are related by volume magnetic susceptibility coefficient.

Although the general form of the $\mathcal{M}^{\alpha\beta}$ and $\mathcal{D}^{\alpha\beta}$ equivalents for the energy-momentum tensors is unknown, one may build two symmetrical energy-momentum tensors, where a division of the stress-energy tensor of electromagnetic field will be obtained, into a tensor representing magnetization-polarization related to charged matter (being the source of the field), and energy-momentum tensor representing electric energy transmission. Next, one may show, that the above leads directly to obtaining the classical equivalent of quantum interpretation seen in QED.

To clarify the above statement one may multiply equation (A6) by μ_r from (3) to get

$$\Upsilon^{\alpha\beta} = -J^{\alpha}\mathbb{A}^{\beta} - \mu_r T^{\alpha\beta} \tag{32}$$

In above

$$-J^{\alpha}\mathbb{A}^{\beta} = \frac{\gamma^2}{(\gamma^2+1)}\frac{B^2}{\mu_o}\cdot\frac{1}{\gamma^2 c^2}U^{\alpha}U^{\beta} \tag{33}$$

what gives first component, describing distribution of magnetic moment. Next, using volume magnetic susceptibility χ from (3) one may introduce the symmetric energy-momentum tensor $\Omega^{\alpha\beta}$ defined as

$$\Omega^{\alpha\beta} \equiv J^{\alpha}\mathbb{A}^{\beta} + \chi T^{\alpha\beta} = -\frac{1}{\mu_o}\mathbb{F}^{\alpha\gamma}\partial_{\gamma}\mathbb{A}^{\beta} \tag{34}$$

where last transformation of the equation comes from (A31), and where according to (A25) above yields

$$-\Upsilon^{\alpha\beta} = \Omega^{\alpha\beta} + T^{\alpha\beta} \tag{35}$$

what yields

$$-\partial_{\beta}\,\Omega^{\alpha\beta} = \partial_{\beta}\,\Upsilon^{\alpha\beta} = f^{\alpha}_{EM}\;;\; -\Omega^{\alpha\beta}\eta_{\alpha\beta} = \varrho c^2 \tag{36}$$

This shows, that $\Omega^{\alpha\beta}$ can be understood as some description of the density of charged matter, experiencing only electromagnetic force. Component $\chi T^{\alpha\beta}$ in equation (34) represents classical relation between Polarization-Magnetization tensor and Electric Displacement tensor, where

$$-\chi T^{00} = u_{B\odot} = \frac{1}{\gamma^2+1}\frac{B^2}{\mu_o} \tag{37}$$

so, by analogy to (27), $\chi T^{\alpha\beta}$ may be also understood as the rank two tensor potential of the electromagnetic field associated with charged matter. All that remains, is to introduce rank two tensor volume magnetic susceptibility $\chi^{\alpha\mu}$ according to the rules of classical electrodynamics

$$\chi^{\alpha\mu} \equiv \tilde{\chi}^{\alpha\mu} + \chi\,\eta^{\alpha\mu} \tag{38}$$

defined in such a way, that

$$\chi^{\alpha\mu}\,T_{\mu}{}^{\beta} = \Omega^{\alpha\beta} \;\;\rightarrow\;\; \tilde{\chi}^{\alpha\mu}\,T_{\mu}{}^{\beta} = J^{\alpha}\mathbb{A}^{\beta} \tag{39}$$

where $\tilde{\chi}^{\alpha\mu}$ seems to be responsible for the self-interaction, resulting in the formation of internal magnetic moments - vortex field associated with elementary particles.

Summarizing,

– $\Omega^{\alpha\beta}$ may be considered as Polarization-Magnetization energy-momentum tensor, describing distribution of charged matter as a sum of rank two tensor electromagnetic potential and energy distribution related to the magnetic moment.

– $T^{\alpha\beta}$ may be considered as Electric Displacement energy-momentum tensor describing electric energy transmission.

Now, one obtains the classical picture (charged matter described by $\Omega^{\alpha\beta}$ exchanging energy of electric field) being equivalent of the description obtained in QED (leptons exchanging bosons), where tensor $\chi^{\alpha\mu}$ from (39) may be further modeled to describe polarization and magnetization by Jones matrices, vectors [26] and symmetry groups [27], analogously as it is done in QED.

In QED picture, according to the final conclusion from [15], if one substitutes (A27) for the current Lagrangian density employed in QED

$$\mathcal{L}_{QED} = \frac{1}{4\mu_o}\mathbb{F}^{\alpha\beta}\,\mathbb{F}_{\alpha\beta} = \frac{1}{2\mu_o}\mathbb{F}^{0\gamma}\partial^{0}\mathbb{A}_{\gamma} = \frac{1}{2}\bar{\psi}(i\hbar c\not{D} - mc^2)\psi \tag{40}$$

one simplifies currently used $\mathcal{L}_{QED}$ and may derive equations that characterize the entire system involving the electromagnetic field where leptons (described by spinors) exchange bosons. However, such a representation of matter can be treated as an equivalent of $\Omega^{\alpha\beta}$ where the description used, reveals only the density of the electromagnetic four-force (36) and four-current (A28).

From this perspective, these equations describe all the forces in the system. As was shown in previous section, four-force densities f^{α}_{gr} and f^{α}_{oth} naturally emerge within the system, and the resultant Lagrangian density duly incorporates this aspect. These forces are now invisible in the equations, because they have been 'absorbed' by the used description of charged matter $\Omega^{\alpha\beta}$, explained in (39) and (36). Above interpretation may thus clarify the challenging quest for identifying quantum gravity as a distinct interaction within Quantum Field Theory. It would also explain the remarkable precision of QED's predictions, provided it indeed characterizes the complete system involving the electromagnetic field.

Since in the system under consideration, all the energy present is the result of the existence of the electromagnetic field (including the energy of charged particles), the above reasoning also leads to the possibility of interpreting the Poynting four-vector $\Upsilon^{0\beta}$ in (32)

$$\Upsilon^{0\beta} = \frac{\gamma^2}{(\gamma^2+1)}\frac{B^2}{\mu_o}\cdot\frac{1}{\gamma c}U^{\beta} - \mu_r T^{0\beta} \tag{41}$$

In the above, the first term is responsible for transferring the energy of magnetic moment what just describes the movement of the density of charged matter. The second element (related to Electric Displacement energy-momentum tensor) can be associated with the transfer of the electric field energy carried by light. From (A27) $T^{00} = \Lambda_{\rho}$, and using (30) it leads to the conclusion that

$$-\mu_r T^{0\beta} = (\, u_{E\odot}\,,\, \overrightarrow{u_{E\odot}}\,) \tag{42}$$

since the energy and momentum densities of photons should be equal, up to the constant c.

This conclusion opens the way to quantum analyzes and makes it possible to provide an equivalent of the presented interpretation for point-like particles.

2.3. Classical and quantum interpretation for point-like particles in flat spacetime

According to interpretation from previous section (42), canonical four-momentum H^{μ} as the volume-integrated $-T^{0\beta}$ (A32) in the description for point-like particles may be associated with a photon, thus

$$cH^{\mu} = (\, H\,,\, c\overrightarrow{p_H}\,) = (\, H\,,\, \overrightarrow{H}\,) \tag{43}$$

which yields $H^{\mu}H_{\mu} = 0$. This confirms the possibility of analyzing light as energy quanta and preserves the fundamental property of equality of energy and momentum of the photon. From (A37) one may also notice, that for a complete description of the behavior of a particle in flat spacetime with an electromagnetic field, it is enough to know Lagrangian and the four-vector $\mathbb{S}^{\mu}$ associated with certain rotation or spin. Unfortunately, $\mathbb{S}^{\mu}$ is unknown, but a quantum solution can be proposed that will shed new light on the interpretation of Quantum Mechanics.

At first, a general method for quantum analysis will be proposed. Referring to conclusions (A24) and (A37), one may introduce new four-vector $\mathcal{E}^{\mu}$ using volume magnetic susceptibility χ from (3) as follows

$$q\mathcal{E}^{\mu} \equiv W^{\mu} - \chi P^{\mu} \tag{44}$$

what yields

$$H^{\mu} - q\mathcal{E}^{\mu} = -q\mathbb{A}^{\mu} \tag{45}$$

Since generalized canonical four-momentum H^{μ} is four-gradient on Hamilton's principal function (A34), therefore, according to freedom of gauge rules, in above equation, four-vector $\mathcal{E}^{\mu}$ is just another gauge of $\mathbb{A}^{\mu}$. Also for any other scalar α, four-vectors $q\mathcal{E}^{\mu} \pm \partial^{\mu}\alpha$ and $q\mathbb{A}^{\mu} \pm \partial^{\mu}\alpha$ always will be an electromagnetic four-potential.

One may thus introduce quantum wave function ψ and wave four-vector K^{μ} related to canonical four-momentum in its simplest form equal to

$$\psi \equiv e^{-iK^{\mu}X_{\mu}}\,;\;\; \hbar K^{\mu} \equiv H^{\mu} \;\;\rightarrow\;\; i\hbar\,\partial^{\mu}\,\psi = H^{\mu}\,\psi \tag{46}$$

Then, acording to (8) and (A24), one may rewrite (45) as just

$$(i\,\hbar\,\partial^{\mu} - q\mathcal{E}^{\mu})\,\psi = \mu_r\,P^{\mu}\,\psi \tag{47}$$

The above equation can be tested in many different quantum applications, which will allow to definitively confirm or deny the validity of the approach proposed in the Alena Tensor theory. An example of such use can be the quantum calculation of the structure of a selected atom. However, in order to be able to use the above formula in calculations, in the following a transformation of the above equation will be presented, so that this equation can be applied to commonly used tools of Quantum Mechanics. In the following steps it will be shown, that the currently used main quantum equations may be actually considered as a result or approximation of the above equation.

At first, using freedom of gauge rules and conclusions (A37) and (A39), one may introduce electromagnetic four-potential $-\Sigma^{\mu}$ defined in following way

$$\Sigma^{\mu} \equiv P^{\mu} + \frac{\varrho c^2 \gamma^2}{p} P^{\beta} + \frac{\varrho c^2}{p} \mathbb{S}^{\mu} + Y^{\mu} \tag{48}$$

what yields

$$H^{\mu} = \Sigma^{\mu} + q\mathbb{A}^{\mu} \;\; \rightarrow \;\; 2H^{\mu} = \Sigma^{\mu} + q\mathcal{E}^{\mu} \tag{49}$$

and thanks to property $H^{\mu}H_{\mu} = 0$ from simple calculations one gets

$$\frac{1}{2}(\Sigma^{\mu}\Sigma_{\mu} + q^2\mathcal{E}^{\mu}\mathcal{E}_{\mu}) = \mu_r^2 m^2 c^2 \tag{50}$$

where according to (3) and (8) one also gets

$$\Sigma^{\mu} = \left(\frac{\mu_r H}{c} \, , \; \overrightarrow{p_H} \; + \; \frac{\chi H}{c^2} \overrightarrow{u} \right) \tag{51}$$

Therefore by introducing vector $\overrightarrow{p_m}$ such that

$$q^2\mathcal{E}^{\mu}\mathcal{E}_{\mu} \equiv \overrightarrow{\Sigma}^{\,2} - \mu_r^2 \overrightarrow{p}_m^{\,2} \tag{52}$$

one may rewrite (50) as just

$$\frac{1}{2}\left(\frac{H^2}{c^2} - \overrightarrow{p}_m^{\,2} \right) = m^2 c^2 \tag{53}$$

what also shows, that with constant H also p_m value is constant. Now, one may perceive the particle using the newly created quantum wave function Ψ, e.g.

$$\Psi \equiv e^{\pm N^{\mu} X_{\mu}} \; ; \;\; \hbar N^{\mu} \equiv \frac{i}{\sqrt{2}} \left(\frac{H}{c} \, , \; \overrightarrow{p_m} \right) \tag{54}$$

to get Klein–Gordon formulation

$$(\hbar^2 \; \Box \; + m^2 c^2)\Psi = 0 \tag{55}$$

One may also consider particle by the corresponding Dirac equation derived in a classical way from above. It would lead to conclusion that the description of a free particle may be considered as contraction of electromagnetic four-potentials with the use of spinor representation.

Finally, it can be shown that the above leads to obtaining the equivalent of the Schrödinger equation. At first it may be noticed, that (A36) and (A37) yield

$$H = mc^2 \left(\gamma + \frac{1}{\gamma} \right) + L = \frac{H^2}{mc^2 \left(\gamma + \frac{1}{\gamma} \right)} - \frac{HL}{mc^2 \left(\gamma + \frac{1}{\gamma} \right)} \tag{56}$$

Next, one may introduce such α, that

$$\partial_t \, \alpha \equiv \frac{H^2}{mc^2 \left(\gamma + \frac{1}{\gamma} \right)} - \mu_r \, H \tag{57}$$

Such α exists, since $\mu_r = \frac{1}{\varepsilon_r}$, where ε_r according to (11) may be expressed by γ and constants, thus $\partial_t \, \alpha$ is function of only γ and constants. Since γ is an integrable function, it can be integrated with respect to t, treating other variables as independent.

Next, referring to (49), (51) and using freedom of gauge rules, a following electromagnetic four-potential $-\widehat{\Sigma}^{\mu}$ can be created

$$\widehat{\Sigma}^{\mu} \equiv \Sigma^{\mu} + \partial^{\mu} \, \alpha = \left(\frac{H^2}{mc^3 \left(\gamma + \frac{1}{\gamma} \right)} \, , \; \overrightarrow{p_H} \; + \; \frac{\chi H}{c^2} \overrightarrow{u} \; - \nabla \;\; \alpha \right) \tag{58}$$

where (56) yields, that second electromagnetic four-potential $q\widehat{\mathbb{A}}^{\mu}$ is

$$q\widehat{\mathbb{A}}^{\mu} \equiv q\mathbb{A}^{\mu} - \partial^{\mu} \, \alpha = \left(\frac{-HL}{mc^3 \left(\gamma + \frac{1}{\gamma} \right)} \, , \; \nabla \;\; \alpha \; - \frac{\chi H}{c^2} \overrightarrow{u} \right) \tag{59}$$

Now using above and (46) one obtains

$$H^{\mu} \, \psi = \widehat{\Sigma}^{\mu} \, \psi + q\widehat{\mathbb{A}}^{\mu} \, \psi \tag{60}$$

what allows to recreate Schrödinger equation by taking zero-components of above four-vectors

$$ic\hbar\,\partial^0\,\psi = -\frac{\hbar^2}{m\left(\gamma+\frac{1}{\gamma}\right)}\nabla^2\quad\psi + cq\widehat{\mathbb{A}}^0\,\psi \tag{61}$$

where $m\left(\gamma+\frac{1}{\gamma}\right)$ may be approximated as 2 m with high accuracy up to velocity $u\approx 0.4c$.

The reasoning presented in this chapter ensures high compliance with the results of Quantum Mechanics and indicates, that the currently used quantum equations for the system with only electromagnetic field may be considered as approximation of the results obtained with the use of Alena Tensor. What is also important, the quantum equations discussed above describe the entire physical system under consideration, including the electromagnetic force, gravity and the Abraham-Lorentz effect, which agrees with the conclusions from the previous chapter. This also means that the non-commutativity of QM is no longer an obstacle to its unification with GR, with the use of Alena Tensor.

In the interpretation presented, one obtains a picture in which gravity and the Abraham-Lorentz effect, in some sense, have always been present in quantum equations. They can be made visible by expanding equation (47) using volume magnetic susceptibility χ from (3), to the form

$$H^\mu = \mu_r\,P^\mu + q\mathcal{E}^\mu = P^\mu + q\mathcal{E}^\mu + \chi\,P^\mu \tag{62}$$

In the classical picture, according to the conclusion (A42) and (6), this leads to the existence of all three forces in the system

$$F^\alpha = U_\beta(\partial^\beta P^\alpha - \partial^\alpha P^\beta) = \frac{1}{\mu_r}\,F^\alpha_{EM} + mc^2\,\partial^\alpha\ln(\mu_r) - \frac{d\ln(\mu_r)}{d\tau}\,P^\alpha \tag{63}$$

where in above, the component $mc^2\,\partial^\alpha\ln(\mu_r) - \frac{d\ln(\mu_r)}{d\tau}\,P^\alpha$ agrees with (5) describing the force related to gravity and the term $\chi\,P^\mu$ in (62) is responsible for the force related to gravity and the Abraham-Lorentz effect.

2.4. Generalization to other fields

To describe uncharged particles related to other fields (e.g. neutrinos), one may also consider generalizing the Alena Tensor to other fields. At this point, however, it seems necessary to introduce a classification of fields that will explain the differences in the approach to their analysis in flat, curved spacetime and in quantum perspective.

Remaining with the previous notation, one may describe the field (e.g. electroweak field) in the system by some generalized field tensor $\mathbb{F}^{\alpha\beta\gamma}$ providing more degrees of freedom, and express Alena Tensor (A1) in flat spacetime as follows

$$T^{\alpha\beta} = \varrho U^\alpha U^\beta - \left(\frac{c^2\varrho}{\Lambda_\rho}+1\right)(\Lambda_\rho\,\eta^{\alpha\beta} - \mathbb{F}^{\alpha\delta\gamma}\,\mathbb{F}^\beta{}_{\delta\gamma}) \tag{64}$$

where

$$\Lambda_\rho \equiv \frac{1}{4}\mathbb{F}^{\alpha\beta\gamma}\mathbb{F}_{\alpha\beta\gamma}\;;\;\xi\,h^{\alpha\beta} \equiv \frac{\mathbb{F}^{\alpha\delta\gamma}\,\mathbb{F}^\beta{}_{\delta\gamma}}{\Lambda_\rho}\;;\;\xi \equiv \frac{4}{\eta_{\alpha\beta}\,h^{\alpha\beta}} \tag{65}$$

The Alena Tensor defined in this way retains most of properties described in appendix, however, it now describes other four-force densities in the system. Total four-force density f^α can be now presented as

$$f^\alpha = \begin{cases} f^\alpha_{fun} \equiv -\partial_\beta\,\mathbb{F}^{\alpha\delta\gamma}\,\mathbb{F}^\beta{}_{\delta\gamma} \quad (\textit{fundamental forces}) \\ + \\ f^\alpha_{gr} \equiv (\eta^{\alpha\beta} - \xi\,h^{\alpha\beta})\partial_\beta\,\varrho c^2 \quad (\textit{related to gravity}) \\ + \\ f^\alpha_{sec} \equiv \frac{\varrho c^2}{\Lambda_\rho} f^\alpha_{fun} \quad (\textit{secondary forces}) \end{cases} \tag{66}$$

Therefore, interactions can be classified based on their properties as:

- fundamental interactions (body forces) related to f^α_{fun}
- related to gravity (against gravity) f^α_{gr}
- secondary interactions (radiation reaction forces) related to f^α_{sec}

where each of above f^α_i four-force density should satisfy the condition $0 = U_\alpha f^\alpha_i$.

Taking into account the conclusions from chapter 2.2, it can be assumed with high probability that the Electroweak Theory describes matter in an analogous way as demonstrated in (35) for electromagnetic interactions, where now $\Upsilon^{\alpha\beta}$ describes the energy-momentum tensor for the electroweak interactions, and $\Omega^{\alpha\beta}$ is still a spinor based description of the matter, this time describing disturbances in the propagation of this field.

This is not so obvious for QCD, due to the strong connection of these interactions with electromagnetism, and it would certainly require further research. However, it seems that the use of Alena Tensor opens up new possibilities in the study of these interactions both in the curvilinear and classical description, as well as in the regime of QFT and QM mathematical apparatus.

3. Discussion

The properties of Alena Tensor seem promising in terms of their further development what could be used in unification theories, in modified theories of gravity [28, 29] or as a method of seeking an explanation for the Dark Sector [29]. For this reason, it is worth discussing the possible use of this tool in selected applications against the background of existing research.

3.1. Dark sector and perspectives for unification of interactions

The first topic discussed will be the issue of the dark sector, for which Alena Tensor brings new interpretation possibilities. Although Dark Energy and Dark Matter are concepts closely related to the General Relativity, their analysis is also carried out e.g. from the perspective of quantum theories and quantum cosmology [30–32].

The use of Alena Tensor indicates that the invariant of the field tensor is responsible for the vacuum energy and the associated cosmological constant [33]. This gives a chance to solve the puzzle of the 'smile of the Cheshire cat' [34] explaining the reason for the appearance of the cosmological constant in Einstein Field equations. Since the first publication of General Relativity, this constant has appeared and disappeared in EFE like Cheshire cat from the book 'Alice's Adventures in Wonderland'. Equation (A13) indicates that cosmological constant is necessary and proposes an explanation of its origin.

Above opens the way to replacing current estimates of the cosmological constant [35] with an attempt to estimate the value of the field tensor invariant. It also becomes possible to search for the expected form of the field tensor based on the experimentally measured value of its invariant, and allows to look for an answer to the question of what fields, apart from the electromagnetic field, should constitute the Alena Tensor.

An example of such an approach seems to be an attempt to estimate the values of magnetic and electric fields based on available background radiation data [36] and an attempt to determine the value of the invariant of the electromagnetic field tensor. Importantly, it also seems that the field invariant in general does not have to be the constant (as a spacetime interval being Lorentz invariant or as the spin). This would give an additional argument for considering the cosmological constant not necessarily as a constant [37, 38], which would be particularly important for solving the Hubble tension problem [39].

Alena Tensor also introduces the possibility of a new interpretation of the forces attributed to Dark Matter. Interpretation presented in section 2.1 drives to conclusion that issues related to Dark Matter may be related to the obtained picture of counter-gravity force (13) or also to e.g. Abraham–Lorentz effect and, in general, does not necessarily involve the existence of additional matter. Modified Alena Tensor may also prove helpful for analysis of Maxwell's equations with axion modifications [40] and attempts to explain Dark Matter based on these particles [41], especially in the context of the results regarding Sigma-8 tension [42]. The connection between gravity and electromagnetism discussed in this article seems also consistent with recent conclusions about the balance of gravity and magnetic fields during the formation of super-massive black holes [43], and perhaps could also help explain the final parsec problem [44] in the description of the black hole collision process.

Analyzing the possible directions of unification of interactions, it can also be noted that the Alena Tensor allows for testing hypotheses regarding the interconnections of fields and the connections of fields with gravity. Fields defined in the way presented in chapter (2.4) allow for quite a lot of freedom in adapting them to the existing division of interactions that emerged in Quantum Mechanics: electroweak, strong and gravitational interactions.

Due to the fundamental importance of electroweak interactions (fermions are the building blocks of matter), it seems that the field strength tensor constituting Alena Tensor should be related to electroweak field, where the rest interactions (strong, gravity and potentially others [45]) could be linked to $f_{gr} + f_{sec}$ four-force density. It would be also supported by conclusions from research on Double Copy Theory [46–48], since it can be assumed that solutions should include perturbative duality between gauge theory and gravity and thus it may be expected, that strong interactions are related to the $f_{gr} + f_{sec}$ four-force density. Also introducing additional fields beyond electromagnetism to Alena Tensor causes an additional spacetime curvature term in Einstein tensor, related to

new fields. Gravity is no longer 'bending of the light path' only, but becomes 'bending of the bosons path of all introduced fields' which, for example, could help to some extent in explaining the strong interactions. Perhaps all this may shed new light on current work on the unification of interactions [49–51].

Finally, when discussing the unification of interactions, it is impossible to ignore the importance of the Higgs field [52]. The adoption of an analysis model based on the Alena Tensor creates new possibilities for relating the geometry of spacetime with a field in general [53] and even based on the simple model presented in A.3, it is possible to analyze relationships between the Higgs field and the electromagnetic field [54, 55]. Additionally, due to the possibility of analyzing the system based on the proposed Lagrangian and generalized canonical four-momentum, it becomes possible to study individual classes of fields in terms of their impact on the phenomenon of symmetry breaking [56, 57].

When building theoretical models, however, one should remember about the limitations related to the adopted analysis method. In curved spacetime, the curvature described by the Einstein tensor will always be related to the four-force densities $f^{\alpha}_{gr} + f^{\alpha}_{sec}$. In flat spacetime, conditions (A23), (A27) and (A32) still seem reasonable.

3.2. Quantum gravity

There is no universal agreement on the approach to developing quantum gravity [58] and so far research is being carried out using different methods in different directions. One of the research directions is canonical quantum gravity [59] with its attempt to quantify the canonical formulation of general relativity, the most promising example of which is Loop Quantum Gravity [60].

Work is also ongoing in the field of string theory, where M-theory [61] seems to be the leading area of research. There are also many other e.g. [62–64] less frequently cited studies that explore different, sometimes unusual [65] research areas.

Against the background of the above research directions, the dualistic approach represented by Alena Tensor seems very promising because it changes the research paradigm in two ways.

The first paradigm shift is that, according to the conclusions presented earlier (and in the appendix), in the description provided by Alena Tensor, the Einstein tensor is not exclusively related to gravity but it also describes phenomena related to four-force density f^{α}_{sec}. This means a change in assumptions and a new way of perceiving prospect of unifying the remaining interactions with gravity.

The second paradigm shift results from the very nature of the dualistic approach and concerns the lack of need to search for quantization methods in curved spacetime. According to the reasoning presented in the article (and in the appendix), if one describes the field in flat spacetime by some field tensor and enters it into the Alena Tensor in the appropriate way, the equations in curved spacetime will naturally turn into the Einstein Field Equations.

The second paradigm shift in particular seems to be extremely important from the point of view of research on quantum gravity phenomena. It also opens new possibilities for studying quantum phenomena in strong gravitational fields.

Current research approaches to quantum problems in strong gravitational field each time require the construction of an appropriate model in which the obtained results can be interpreted, either through careful selection of the observer [66], or making direct use of the principle of equivalence [67], or own, specific approach [68]. It also needed consideration of the specific quantum phenomena occurring in the vicinity of very massive objects, such as the Unruh effect [69] or Hawking radiation [70]. Thanks to the dualistic approach, such research can now be conducted in flat spacetime with fields and then the results can be analyzed in curved spacetime.

One of the natural directions of research seems to be the development of a field tensor that, in curved spacetime, provides the known metrics [71] used to describe gravity, extended by the term related to secondary interactions. The development of such a field tensor seems to be the first step towards building quantum gravity, this time - contrary to the direction described in the previous chapter - from the side of the General Relativity.

Interestingly, because the use of the Alena Tensor indicates the possibility of shaping the metric tensor of spacetime using a field, it also sheds new light on research on new drives [72], including the quantum effects [73] needed to analyze them. Although many QM and QFT problems seem unsolvable [74, 75] using current paradigms, such as the Planck scale problems [76], previously mentioned paradigm shifts may change this situation.

It also seems interesting to search for solutions to the problem of quantization of interactions related to the tensor (A9) in various spacetimes, thus the problem of quantization should be addressed.

3.3. Quantization

To get a full picture of the applicability of the approach based on Alena Tensor, one may consider an example of its application to gravity quantization.

One may start with a choice of proper representation of a metric $g^{\alpha\beta}$ so that the interpretation of time in first quantization will be 'natural'. By 'natural interpretation' of time, it is understood the approach in which, after

the first quantization of Hamiltonian, one gets a proper definition of the time evolution operator in the 'Schrödinger representation', in such a way that

$$U(t,\ t_0) = e^{-iH\cdot(t-t_0)/\hbar} \tag{67}$$

fulfill classical conditions [77]

$$U^{\dagger}(t,\ t_0)U(t,\ t_0) = I \tag{68}$$

$$|\psi(t_0)\rangle = U(t_0,\ t_0)|\psi(t_0)\rangle \tag{69}$$

$$U(t,\ t_0) = U(t,\ t_1)U(t_1,\ t_0) \tag{70}$$

This means that, in general, it should be possible to incorporate the Lagrangian formalism for the Gauge fields. Therefore, for the field strength tensor

$$F^a_{\mu\nu} = \partial_\mu A^a_\nu - \partial_\nu A^a_\mu + g f^{abc} A^b_\mu A^c_\nu \tag{71}$$

one needs to define proper commutator

$$[t_a,\ t_b] = i f^{abc} t_c \tag{72}$$

As it was show in [78] this can be done by rewriting $g^{\alpha\beta}$ in the $(3+1)$-split in Geroch decomposition manner. This approach solves the proper initial value problem, since now spacetime can be interpreted as the evolution of space in time, with interpretation of time that is consistent with Quantum Mechanics: time as a distinguished, absolute, external, global parameter. A summary of full formalism has been presented many times, last and the modern one can be found in [79], where computation rules look as follows

$$\{\gamma_{ij},\ \pi^{kl}\} = \frac{1}{2}(\delta^k_i\delta^l_j + \delta^k_j\delta^l_i)\delta^{(3)}(x-y) \tag{73}$$

The above approach makes it possible to introduce gravity into Quantum Mechanics in form of canonical quantization and couple this field with other interactions in regular manner. In such picture gravity acts as just another quantum field that could be incorporated into the Standard Model Lagrangian and interact with other fields on the same principles. The only difference is that we are bound to only one representation of the metric $g^{\alpha\beta}$ with $(3+1)$-split Geroch decomposition. However, it may be transformed to other, more convenient coordinate systems when quantum phenomena can be negligible.

Presented approach opens a natural way to implement representation of tensor $g^{\alpha\beta}$ into the Alena Tensor (A1) for better understanding the interpretation of GR in the big scale. It also opens the possibility to look for quantum gravity phenomena on the small scale, where perturbation approach as quantum and gravity interaction are on the same level of magnitude. The most promising application of this approach could be implementing this calculations to Hawking radiation phenomena on the Planck scales, as the original calculations are questioned by other authors [80, 81].

New observation methods allow to look for a quantum gravity phenomenon in the present or near future data that could test the boundaries of GR in the classical approach. One of the most promising directions is the rise of gravitational wave (GW) astronomy. It might be worth investigating the post-merge echoes that occur because of the stimulated emission of Hawking radiation after compact binary merger events involving stellar black holes. This could be a promising way to search for deviations from General Relativity and could serve as evidence for the quantum structure of black hole horizons. Present methods used to model this phenomenon in modified theories of gravity are extremely challenging in Numerical Relativity and could provide inconclusive observation interpretations [82]. The approach presented in this paper may also help obtain results without using effective model echoes within the framework of linear perturbation theory.

4. Conclusion

As presented in the article (and in the appendix), the possibility of using a new tool, Alena Tensor, seems to open up new research possibilities both in terms of searching for the relationship between QFT and GR [83], as well as in terms of connections between many phenomena previously analyzed separately: in quantum or classical description, curvilinear or in flat spacetime, or, for example, the possibility of combining the interpretation of fluid dynamics with field theory. Such an analysis may prove particularly interesting in the context of cosmology and the study of quantum phenomena in the early universe [84]. The lines of unification proposed by Alena Tensor can be visualized as in figure 2 below.

By appropriately selecting field tensors and testing hypotheses regarding their relationship with the Einstein tensor in curved spacetime, it is possible to search for new interpretations for Dark Matter, as well as to analyze the relationships of the invariants of these field tensors with the cosmological constant. By adopting a new interpretation of the cosmological constant as an invariant of the field tensor, possibilities also open up to



Figure 2. Connections between knowledge areas, introduced by Alena Tensor. Source: self-made illustration created with the assistance of AI.

explain contradictory experimental data for cosmological phenomena, because the field tensor invariant, in general, does not have to be constant.

Due to the high flexibility of the Alena Tensor in the selection of fields, it also seems to be a good tool for testing hypotheses regarding the unification of interactions. Such research can be conducted in the regime of the QFT mathematical apparatus and, importantly, thanks to a clear interpretation of the four-divergence of the field stress-energy tensor (four-force density), obtained results would also lead to obtaining an interpretation of quantum interactions in the classical description. It could be a major milestone in combining known QFT results with the classical description of interactions.

Finally, one can also seek a quantum description of gravity in new ways, taking advantage of the paradigm shifts that Alena Tensor brings with it. This does not mean that the problems associated with quantizing fields in curved spacetime disappear and the behavior of quantum fields, when changing the metric tensor, will still require careful analysis. However, it seems that thanks to the dualistic description provided by Alena Tensor, these analysis may be much easier.

The Alena Tensor can be proven based on derived quantum equations (47) or (61), which opens the way to new experimental and theoretical research. Further research on Alena Tensor may also lead to its important transformations and generalizations, as well as to the design of experiments in terms of the sought properties that match the experimental data. And all this has a chance to bring us one step closer to the next image that will connect the previously scattered puzzles of knowledge.

5. Statements

All data that support the findings of this study are included within the article (and any supplementary files).

During the preparation of this work the authors did not use generative AI or AI-assisted technologies, except for generating the elements of figure 2.

Authors did not receive support from any organization for the submitted work.

Authors have no relevant financial or non-financial interests to disclose.

Both authors contributed equally.

Data availability statement

All data that support the findings of this study are included within the article (and any supplementary files).

Appendix. Summary of conclusions from previous publications about Alena Tensor

A.1. Alena Tensor and main definitions

This appendix summarizes the state of knowledge about Alena Tensor based on recent publications and systematizes existing conclusions in the context of further applications.

Alena Tensor is the central object of the method described in [14, 15]. It is a stress-energy tensor, which can be interpreted in flat and curved spacetime. The Alena Tensor $T^{\alpha\beta}$ has the following form

$$T^{\alpha\beta} = \varrho\ U^{\alpha}U^{\beta} - (c^2\varrho + \Lambda_{\rho})(g^{\alpha\beta} - \xi\ h^{\alpha\beta}) \tag{A1}$$

Designations used:

- $g^{\alpha\beta}$ is a metric tensor by which the spacetime of a physical system is considered,
- $1/\xi \equiv \frac{1}{4}\, g_{\mu\nu}\ h^{\mu\nu}$,
- $\varrho \equiv \varrho_o\gamma$ where ϱ_o is rest mass density and γ is Lorentz gamma factor,
- $\varrho\ U^{\alpha}$ is four-momentum density in the system, in accordance with the postulate raised in the description to equation (11) from publication [14],
- $h^{\alpha\beta}$ is the metric tensor of curved spacetime in which all motion takes place along geodesics and it is related to the field tensor, which will be explained next,
- Λ_{ρ} is related to the invariant of the field tensor, which will be explained next.

The field present in the system is described by some field tensor, e.g. $\mathbb{F}^{\alpha\beta\gamma}$, which may be widely configured. To simplify the reasoning, it will be assumed that field is described by $\mathbb{F}^{\beta\gamma}$ representing electromagnetic field, but the properties described here are general and apply to the field in a broader sense.

Considering a system with only an electromagnetic field, where this field is described by a tensor $\mathbb{F}^{\beta\gamma}$ one gets the following relationships

$$h^{\alpha\beta} \equiv 2\ \frac{\mathbb{F}^{\alpha\delta}\, g_{\delta\gamma}\, \mathbb{F}^{\beta\gamma}}{\sqrt{\mathbb{F}^{\alpha\delta}\, g_{\delta\gamma}\, \mathbb{F}^{\beta\gamma}\, g_{\mu\beta}\, \mathbb{F}_{\alpha\eta}\, g^{\eta\xi}\, \mathbb{F}^{\mu}{}_{\xi}}} \tag{A2}$$

which provides the property $h^{\alpha\beta}\, g_{\mu\beta}\, h_{\alpha}{}^{\mu} = 4$, and

$$\Lambda_{\rho} \equiv \frac{1}{4\mu_o}\mathbb{F}^{\alpha\mu}\, g_{\mu\gamma}\, \mathbb{F}^{\beta\gamma} g_{\alpha\beta} \tag{A3}$$

where μ_o is vacuum magnetic permeability. Considered system without a field vanishes (for example, without EM field there is no light or charged particles, and other fields and particles associated with them are absent if a system containing only EM field is considered). The dependence of $h^{\alpha\beta}$ on the field tensor (A2), indicates that spacetime in this approach should be actually understood as some method to perceive the field (indeed, in a curvilinear description, geodesics describe the paths of light propagation in considered system).

The stress–energy tensor for electromagnetic filed, denoted as $\Upsilon^{\alpha\beta}$ may be thus presented in a way that relates the field to the metric tensor of curved spacetime

$$\Upsilon^{\alpha\beta} \equiv \Lambda_{\rho}(g^{\alpha\beta} - \xi\ h^{\alpha\beta}) = \Lambda_{\rho} g^{\alpha\beta} - \frac{1}{\mu_o}\mathbb{F}^{\alpha\delta}\, g_{\delta\gamma}\, \mathbb{F}^{\beta\gamma} \tag{A4}$$

This connection of the field with the $h^{\alpha\beta}$ tensor opens up wide possibilities of unification, discussed in the main article.

The pressure p in the system is equal to

$$p \equiv c^2\varrho + \Lambda_{\rho} \tag{A5}$$

where it was shown in [15] that p is negative. This allows (A1) to be written as

$$T^{\alpha\beta} = \varrho\ U^{\alpha}U^{\beta} - \frac{p}{\Lambda_{\rho}}\Upsilon^{\alpha\beta} \tag{A6}$$

The remaining tensors that describe the system are defined as follows

$$R^{\alpha\beta} \equiv 2\varrho\ U^{\alpha}U^{\beta} - p\ g^{\alpha\beta} \tag{A7}$$

its trace R

$$R \equiv R^{\alpha\beta}\ g_{\alpha\beta} = -2p - 2\Lambda_{\rho} \tag{A8}$$

and tensor $G^{\alpha\beta}$ as

$$G^{\alpha\beta} \equiv R^{\alpha\beta} - \frac{1}{2}R\ \xi\ h^{\alpha\beta} \tag{A9}$$

which allows to rewrite (A1) as

$$G^{\alpha\beta} - \Lambda_{\rho}\ g^{\alpha\beta} = 2\ T^{\alpha\beta} + \varrho c^{2}(g^{\alpha\beta} - \xi\ h^{\alpha\beta}) \tag{A10}$$

The above definitions allow to consider flat spacetime, curved spacetime, and all intermediate states, in which spacetime is partially curved and part of the motion results from the existence of residual fields. One may analyze boundary solutions: flat spacetime with fields and curved spacetime without fields.

A.2. Behavior of the system in curved spacetime

Considering $g^{\alpha\beta}$ as equal to $h^{\alpha\beta}$ one obtains that it yields $\xi = 1$, therefore the whole part of Alena Tensor related to fields vanishes. It yields

$$T_{\alpha\beta} = \varrho\ U_{\alpha}U_{\beta} \tag{A11}$$

The value of tensor $G_{\alpha\beta}$ becomes

$$G_{\alpha\beta} = R_{\alpha\beta} - \frac{1}{2}R\ h_{\alpha\beta} \tag{A12}$$

and (A10) reduces to

$$G_{\alpha\beta} - \Lambda_{\rho}\ h_{\alpha\beta} = 2\ T_{\alpha\beta} \tag{A13}$$

Therefore, in curved spacetime, $R_{\alpha\beta}$ acts as Ricci tensor and $G_{\alpha\beta}$ acts as Einstein curvature tensor, both with an accuracy of $\frac{4\pi G}{c^{4}}$ constant, where cosmological constant Λ is related to the invariant of the field tensor

$$\Lambda = -\frac{4\pi G}{c^{4}}\Lambda_{\rho} \tag{A14}$$

where Λ_{ρ} is negative. Equation (A13) can be further analyzed using known tools for considering metrics in General Relativity, taking into account the knowledge of the field tensor used to build the Alena Tensor.

Since covariant four-divergences of $T_{\alpha\beta}$ and $G_{\alpha\beta}$ vanish, therefore they represent curvature tensors, related to corresponding four-force densities present in flat Minkowski spacetime. For this reason it is worth to consider behavior of the system in flat Minkowski spacetime.

A.3. Behavior of the system in flat Minkowski spacetime

Considering $g^{\alpha\beta}$ as equal to $\eta^{\alpha\beta}$ Minkowski metric tensor, thanks to the amendment to the Continuum Mechanics explained in equations (13) - (21) of publication [14]

$$\partial_{\alpha}U^{\alpha} = -\frac{d\gamma}{dt} \quad \rightarrow \quad \partial_{\alpha}\ \varrho\ U^{\alpha} = 0 \tag{A15}$$

total four-force density f^{α} acting in the system is equal to

$$f^{\alpha} \equiv \partial_{\beta}\ \varrho\ U^{\alpha}U^{\beta} \tag{A16}$$

and for considered system, it is the sum of electromagnetic (f^{α}_{EM}), related to gravity (f^{α}_{gr}) and other (f^{α}_{oth}) four-force densities, where

$$f^{\alpha} = \begin{cases} f^{\alpha}_{EM} \equiv \partial_{\beta}\ \Upsilon^{\alpha\beta} \quad (electromagnetic) \\ + \\ f^{\alpha}_{gr} \equiv (\eta^{\alpha\beta} - \xi\ h^{\alpha\beta})\partial_{\beta}\ \varrho c^{2} \quad (related\ to\ gravity) \\ + \\ f^{\alpha}_{oth} \equiv \frac{\varrho c^{2}}{\Lambda_{\rho}} f^{\alpha}_{EM} \quad (other) \end{cases} \tag{A17}$$

In above, f^{α}_{gr} is not an interaction between bodies, but appears to result from the bending of the direction of electromagnetic field energy transport by the energy density gradient. The meaning of the four-force density f^{α}_{gr} is discussed in the main article and it is not so much the force of gravity as the force resulting from opposing

gravity. Equation (A17) yields

$$\partial_\beta T^{\alpha\beta} = 0 \tag{A18}$$

and

$$\partial_\beta G^{\alpha\beta} = f^\alpha_{gr} + f^\alpha_{oth} \tag{A19}$$

The above result shows, that when using the Alena Tensor, it should be assumed that the Einstein tensor does not describe the curvature associated with gravity alone.

Neglecting other forces (as we currently do in known solutions for GR), one actually approximately obtains metric tensors responsible for gravity alone. However, the total value of the Einstein tensor corresponds to the curvature associated with the density of the four-forces from equation (A19). This means that the above approach can be used to search for the causes of disturbances between observations and the expected motion resulting from gravitational equations, which is currently attributed entirely to Dark Matter [85].

The meaning of the four-force density f^α_{oth} is discussed in the main article.

One may also introduce an additional tensor $\Pi^{\alpha\beta}$ which turns out to play a role of deviatoric stress tensor [86]

$$\Pi^{\alpha\beta} \equiv -c^2 \varrho\ \xi\ h^{\alpha\beta} \tag{A20}$$

To demonstrate this, Alena Tensor can be represented in flat Minkowski spacetime as

$$T^{\alpha\beta} = \varrho\ U^\alpha U^\beta - p\ \eta^{\alpha\beta} - \Pi^{\alpha\beta} + \Lambda_\rho \xi\ h^{\alpha\beta} \tag{A21}$$

Now, vanishing four-divergence of the above

$$f^\alpha = \partial^\alpha p + \partial_\beta \Pi^{\alpha\beta} + f^\alpha_{EM} \tag{A22}$$

express relativistic equivalence of Cauchy momentum equation (convective form) [87]. The above representation therefore allows for the analysis of the system using the tools of continuum mechanics. From this perspective, f_{EM} appears as a body force, while the remaining forces are the effect of fluid dynamics [88] and could be modeled e.g. with help of Navier–Stokes equations [89, 90].

By imposing following condition on normalized Alena Tensor as described in [15]

$$0 = \partial_\beta \left(\frac{T^{\alpha\beta}}{\eta_{\mu\gamma} T^{\mu\gamma}} \right) + \partial^\alpha \ln\left(\eta_{\mu\gamma} T^{\mu\gamma}\right) \tag{A23}$$

one obtains further simplification. Some gauge of electromagnetic four-potential denoted as $\mathbb{A}^\mu$ may be expressed as

$$\mathbb{A}^\mu \equiv -\frac{\Lambda_\rho}{p}\frac{\varrho_o}{\rho_o} U^\mu \tag{A24}$$

where ρ_o denotes rest charge density in the system. It also simplifies Alena Tensor in flat Minkowski spacetime to

$$T^{\alpha\beta} = \frac{1}{\mu_o}\mathbb{F}^{\alpha\gamma}\partial^\beta \mathbb{A}_\gamma - \Lambda_\rho \eta^{\alpha\beta} \tag{A25}$$

and leads to the explicit form of gravitational four-force density

$$f^\alpha_{gr} = \varrho\left(\frac{d\ \ln(p)}{d\tau} U^\mu - c^2 \partial^\mu \ln(p) \right) \tag{A26}$$

Both Lagrangian density ($\mathcal{L}$) and Hamiltonian density ($\mathcal{H} = T^{00}$) for the system appear to be related to invariant of the field tensor

$$\mathcal{L} = \mathcal{H} = \Lambda_\rho \tag{A27}$$

where it was shown in [15] that

$$\frac{\partial \Lambda_\rho}{\partial \mathbb{A}_\alpha} = \partial_\nu \left(\frac{\partial \Lambda_\rho}{\partial(\partial_\nu \mathbb{A}_\alpha)} \right) = -J^\alpha \tag{A28}$$

In above

$$J^\alpha = \rho_o \gamma U^\alpha \tag{A29}$$

is electric four-current and according to (A15) its four-divergence vanishes. Equation (A27) indicates, that in this solution there is no potential in the classical sense and dynamics of the system depends on itself. This is a clear analogy to main GR equation and something that should be expected from a GR-equivalent description of the system in flat spacetime.

Finally, electromagnetic field energy density Υ^{00} was calculated in equation (57) of [15] as

$$\Upsilon^{00} = \frac{\Lambda_\rho}{p}(\varrho c^2\gamma^2 - \Lambda_\rho) \tag{A30}$$

and second representation of the stress-energy tensor was shown in equation (38) of [15] as

$$T^{\alpha\beta} = \frac{p}{\varrho c^2}\,\partial_\gamma \frac{1}{\mu_o}\mathbb{F}^{\alpha\gamma}\mathbb{A}^\beta \tag{A31}$$

A.4. Dynamics of point-like particles in flat spacetime

It was also shown in [15], that

$$H^\beta \equiv \left(\frac{H}{c}\,,\,\overrightarrow{p_H}\right) \equiv -\frac{1}{c}\int T^{0\beta}\,d^3x \tag{A32}$$

in flat spacetime acts as canonical four-momentum for the point-like particle, and for the system with electromagnetic field, four-divergence of H^β vanishes due to the Poynting theorem. Hamiltonian for point-like particle is thus

$$H = -\int \Lambda_\rho\,d^3x \tag{A33}$$

The action S (Hamilton's principal function) for the point-like particle was derived in [15] as

$$-S = H^\beta X_\beta = mc^2\tau + \int p\,d^4x = P^\beta X_\beta - mc^2\tau \tag{A34}$$

where P^β is four-momentum and τ is particle's proper-time. One may denote in the above equation Pressure-Volume work (pressure potential energy) as W_{pv}

$$W_{pv} \equiv -\int p\,d^3x \tag{A35}$$

and it has positive value. Denoting F^β as total four-force acting on the particle one may notice that Lagrangian L for the particle may be understood as the Lagrangian for a particle of some perfect fluid [91]

$$-L = \frac{1}{\gamma}F^\beta X_\beta = \frac{mc^2}{\gamma} - W_{pv} \tag{A36}$$

thus it may be also analyzed from the perspective of the laws of Thermodynamics.

Hamilton's principal function (action) (A34) vanishes for the inertial system. It clearly shows that inertial systems in this approach do not exist and should be considered as some abstract idealization.

According to [15], mentioned canonical four-momentum may be expressed as

$$H^\mu = P^\mu + W^\mu = -\frac{\gamma L}{c^2}U^\mu + \mathbb{S}^\mu \tag{A37}$$

where L is Lagrangian for point-like particle, $\mathbb{S}^\mu$ due to its property $\mathbb{S}^\mu U_\mu = 0$, seems to be some description of rotation or spin, and where W^μ describes the transport of energy due to the field. It can be expressed in a generalized way as

$$W^\mu = X_\beta \partial^\mu P^\beta - \partial^\mu mc^2\tau \tag{A38}$$

For considered system with electromagnetic field it was calculated in [15] as

$$W^\mu = q\mathbb{A}^\mu + \frac{\varrho c^2\gamma^2}{p}P^\beta + \frac{\varrho c^2}{p}\mathbb{S}^\mu + Y^\mu \tag{A39}$$

where Y^μ is the volume integral of the Poyinting four-vector

$$Y^\beta = \int \Upsilon^{0\beta}\,d^3x \tag{A40}$$

and

$$\mathbb{S}^\beta = \int \frac{\varepsilon_o\Lambda_\rho}{\gamma c\rho_o}\,\mathbb{F}^{0\mu}\partial_\mu U^\beta\,d^3x \tag{A41}$$

where ε_o is electric vacuum permittivity.

Since in (A37) W^μ is just 'other gauge' of $-P^\mu$ thus in the classical description for such a system occurs

$$F^\alpha = U_\beta(\partial^\beta P^\alpha - \partial^\alpha P^\beta) = U_\beta(\partial^\alpha W^\beta - \partial^\beta W^\alpha) \tag{A42}$$

where $U_\beta\partial^\alpha P^\beta = 0$ vanishes, due to the property of Minkowski metric $\partial^\alpha U_\beta U^\beta = 0$.

ORCID iDs

Piotr Ogonowski https://orcid.org/0000-0001-5526-3439

References

[1] Böhmer C G and Downes R J 2014 *International Journal of Modern Physics* D **23** 1442015
[2] Afonso V I, Olmo G J, Orazi E and Rubiera-Garcìa D 2018 The European Physical Journal. C *Particles and Fields* **78** 866
[3] Shalyt-Margolin A 2021 *International Journal of Theoretical Physics* **60** 1858–69
[4] Abdalla E and Marins A 2020 *International Journal of Modern Physics* D **29** 2030014
[5] Goenner H 2004 *Living Rev. Relativ.* **7**
[6] Halverson J and Sung B 2022 *Physical Review* D **105** 126012
[7] Canepa A 2019 *Reviews in Physics* **4** 100033
[8] Rowlands P 2021 An approach to grand unification *Journal of Physics: Conference Series* vol 2081 (IOP Publishing) 012010
[9] Mahanta M 1984 *Annalen der Physik* **496** 357–71
[10] Sanchez N G 2019 *International Journal of Modern Physics* D **28** 1950055
[11] Grimmer D 2023 Introducing the ise methodology: A powerful new tool for topological redescription arXiv:2303.04130
[12] Beach R J 2024 *Annals of Physics* **465** 169661
[13] Torromé R G 2020 *International Journal of Geometric Methods in Modern Physics* **17** 2050060
[14] Ogonowski P 2023 *International Journal of Modern Physics* D **2350010** 15
[15] Ogonowski P 2023 *Frontiers in Physics* **11** 1264925
[16] Polonyi J 2019 *International Journal of Modern Physics* A **34** 1950077
[17] Roberts B L and Marciano W J 2010 *Lepton Dipole Moments* 20 (World Scientific)
[18] Masood S and Mein H 2019 arXiv:1901.08569
[19] Wen M, Keitel C H and Bauke H 2017 *Physical Review* A **95** 042102
[20] Wen M, Bauke H and Keitel C H 2016 *Scientific reports* **6** 31624
[21] d'Inverno R and Vickers J 2022 *Introducing Einsteinaposs Relativity: A Deeper Understanding* (Oxford University Press)
[22] Herrero-Valea M, Liberati S and Santos-Garcia R 2021 *Journal of High Energy Physics* **2021** 1–32
[23] Schleich K and Witt D 2008 *Canadian Journal of Physics* **86** 591–5
[24] Ikeda T N and Sato M 2019 *Physical Review B* **100** 214424
[25] Gonano C A, Zich R E and Mussetta M 2015 *Progress In Electromagnetics Research B* **64** 83–101
[26] Karlsson M 2013 The connection between polarization calculus and four-dimensional rotations arXiv:1303.1836
[27] Başkal S and Kim Y 2018 Lorentz group in ray and polarization optics *Mathematical Optics* (CRC Press) 303–40
[28] Böhmer C G and Jensko E 2024 *Fortschritte der Physik* 2300216
[29] Capozziello S, Mazumdar A and Meluccio G 2024 *Physics of the Dark Universe* **45** 101517
[30] Foster J W, Kumar S, Safdi B R and Soreq Y 2022 *Journal of High Energy Physics* **2022**
[31] Das S, Sharma M K and Sur S 2021 *Physical Sciences Forum* **2** 55
[32] Ng Y 2016 *Journal of Physics: Conference Series* **845** 012001
[33] Peebles P J E and Ratra B 2002 *Reviews of Modern Physics* **75** 559–606
[34] Das Gupta P 2012 *Resonance* **17** 254–73
[35] Carlip S 2019 *Physical Review Letters* **123** 131302
[36] Lewis A 2003 *Physical Review* D **68** 083509
[37] Shapiro I L, Sola J, Espana-Bonet C and Ruiz-Lapuente P 2003 *Physics Letters* B **574** 149–55
[38] Dvali G and Vilenkin A 2001 *Physical Review* D **64** 063509
[39] Di Valentino E, Mena O, Pan S, Visinelli L, Yang W, Melchiorri A, Mota D F, Riess A G and Silk J 2021 *Classical and Quantum Gravity* **38** 153001
[40] Li T, Zhang R J and Dai C 2023 *Journal of High Energy Physics* **2022** 1–18
[41] Lee Y, Yang B, Yoon H, Ahn M, Park H B, Min B, Kim D and Yoo J 2022 *Physical Review Letters* **128** 241805
[42] Joseph M, Aloni D, Schmaltz M, Sivarajan E N and Weiner N 2023 *Physical Review* D **108** 023520
[43] Zou F, Yu Z, Brandt W, Tak H, Yang G and Ni Q 2024 *The Astrophysical Journal* **964** 183
[44] Milosavljević M and Merritt D 2003 The final parsec problem *AIP Conference Proceedings* vol 686 (American Institute of Physics) 201–10
[45] 2022 *Nature Physics* **18** 277–82
[46] Dìaz-Jaramillo F, Hohm O and Plefka J 2021 *Physical Review* D **105** 045012
[47] Spallucci E and Smailagic A 2022 *Physics Letters* B **831** 137188
[48] Easson D A, Manton T and Svesko A 2021 *Physical Review Letters* **127** 271101
[49] Guangzhou G 2019 *Physics Essays* **32** 399–410
[50] Davighi J and Tooby-Smith J 2022 *Journal of High Energy Physics* **2022** 193
[51] Sarfatti J 2010 *Bulletin of the American Physical Society*
[52] Lisi A, Smolin L and Speziale S 2010 *Journal of Physics A: Mathematical and Theoretical* **43** 445401
[53] Madore J 2008 *International Journal of Geometric Methods in Modern Physics* **05** 265–9
[54] Damgaard P H and Heller U M 1988 *Nuclear Physics* **309** 625–54
[55] Nielsen N 2014 *Physical Review* D **90** 016010
[56] Leder E 2020 *Symmetry, Symmetry Breaking, and the Current View of the Dirac Monopole* Portland State University (https://doi.org/10.15760/etd.7261)
[57] Quigg C 2007 *Reports on Progress in Physics* **70** 1019–53
[58] Krasnov K and Percacci R 2018 *Classical and Quantum Gravity* **35** 143001
[59] Thiemann T 2020 *Frontiers in Physics* **8** 548232
[60] Ashtekar A and Bianchi E 2021 *Reports on Progress in Physics* **84** 042001
[61] Albertini F, Del Zotto M, Garcìa Etxebarria I and Hosseini S S 2020 *Journal of High Energy Physics* **2020** 1–46
[62] Loll R 2019 *Classical and Quantum Gravity* **37** 013002
[63] Fernandes P G, Carrilho P, Clifton T and Mulryne D J 2022 *Classical and Quantum Gravity* **39** 063001

[64] Wani S S, Quach J Q and Faizal M 2021 *Europhysics Letters* **139** 62002
[65] Epstein H I 2021 *Physics Essays* **34** 429–63
[66] Augousti A, Gawełczyk M, Siwek A and Radosz A 2011 *European Journal of Physics* **33** 1
[67] Demir D 2022 *Physical Review* A **106** 022215
[68] Pailas T 2020 *Quantum Reports* **2** 414–41
[69] Chen A 2023 *Physical Review* D **107** 056014
[70] Kolobov V I, Golubkov K, de Nova J R M and Steinhauer J 2021 *Nature Physics* **1** 6
[71] Carroll S M 2019 *Spacetime and Geometry* (Cambridge University Press)
[72] Alcubierre M and Lobo F S 2017 Warp Drive Basics *Wormholes, Warp Drives and Energy Conditions* (Springer Cham) 257–79
[73] Lundblad N *et al* 2023 *Quantum Science and Technology* **8** 024003
[74] Tachikawa Y 2023 *International Journal of Theoretical Physics* **62** 1–13
[75] Noce C and Romano A 2022 *Encyclopedia* **2** 1517–27
[76] Diósi L 2019 *Journal of Physics: Conference Series* **1275**
[77] Sakurai J and Napolitano J 2017 *Modern Quantum Mechanics* (Cambridge University Press) ISBN 9781108422413
[78] Ogonowski P and Skindzier P 2013 Maxwell-like picture of general relativity and its planck limit arXiv:1301.2758
[79] Bergstedt V 2020 *Spacetime as a Hamiltonian Orbit and Geroch's Theorem on the Existence of Fermions* Uppsala University, Disciplinary Domain of Science and Technology, Physics, Department of Physics and Astronomy
[80] Helfer A D 2003 *Reports on Progress in Physics* **66** 943–1008
[81] Brout R, Massar S, Parentani R and Spindel P 1995 *Physical Review* D **52** 4559–68
[82] Abedi J, Longo Micchi L F and Afshordi N 2023 *Phys. Rev.* D **108** 044047
[83] Yang R Q, Liu H, Zhu S, Luo L and Cai R G 2020 *Physical Review Research* **2** 023107
[84] Carloni S, Cianci R, Corradini O, Flachi A, Vignolo S and Vitagliano V 2023 Avenues of quantum field theory in curved spacetime, genova, 14-16 sep 2022 *Journal of Physics: Conference Series* vol 2531 (IOP Publishing) 011001
[85] Bertone G and Hooper D 2018 *Reviews of Modern Physics* **90** 045002
[86] Surana K S, Joy A D, Kedari S R, Nuñez D E, Reddy J and Wongwises S 2017 A nonlinear constitutive theory for deviatoric cauchy stress tensor for incrompressible viscous fluids **3** 1221–40 https://hdl.handle.net/1808/27331
[87] Goraj R 2016 *Applied Mathematics-a Journal of Chinese Universities Series B* **07** 1068–73
[88] Romatschke P and Romatschke U 2019 *Relativistic Fluid Dynamics In and Out of Equilibrium: and Applications to Relativistic Nuclear Collisions* (Cambridge University Press)
[89] Bredberg I, Keeler C, Lysov V and Strominger A 2012 *Journal of High Energy Physics* **2012** 1–18
[90] Lasukov V 2019 *Russian Physics Journal* **62** 778–93
[91] Manoff S 2003 Lagrangian theory for perfect fluids arXiv:gr-qc/0303015